

Table 2. Number, incidence rate ¹, median days away from work ² and relative standard errors ³ of occupational injuries and illnesses involving days away from work ⁴ to selected parts of body with musculoskeletal disorders⁵ in private industry for Georgia, 2007

Part of body affected	Total Cases	Incidence Rate	Median Days	Relative Standard Error
All Selected Parts	6,310	21.5	8	4.7
1 Neck- Including Throat	190	0.7	6	13.9
10 Neck- except internal location of diseases or disorders	190	0.7	6	13.9
2 Trunk	4,560	15.5	8	4.9
21 Shoulder- including clavicle- scapula	770	2.6	21	7.8
22 Chest- including ribs- internal organs	40	0.1	3	31.2
220 Chest- except internal location of diseases or disorders	40	0.1	3	31.2
23 Back- including spine- spinal cord	3,100	10.5	6	5.3
230 Back- including spine- spinal cord- unspecified	1,210	4.1	6	6.7
231 Lumbar region	1,610	5.5	6	6.2
232 Thoracic region	230	0.8	2	12.8
238 Multiple back regions	40	0.1	7	29.5
24 Abdomen	350	1.2	15	10.7
240 Abdomen- except internal location of diseases or disorders	50	0.2	7	26.4
241 Internal abdominal location- unspecified	200	0.7	15	13.7
245 Intestines- peritoneum	100	0.3	32	18.9
2450 Intestines- peritoneum- unspecified	100	0.3	32	18.9
25 Pelvic region	210	0.7	5	13.4
254 Groin	200	0.7	5	13.6
28 Multiple trunk locations	80	0.3	16	20.6
3 Upper extremities	810	2.7	9	7.7
31 Arm(s)	300	1.0	4	11.5
310 Arm(s)- unspecified	140	0.5	2	16.2
311 Upper arm(s)	20	0.1	60	38.5
312 Elbow(s)	90	0.3	4	20.0
313 Forearm(s)	30	0.1	14	33.6
32 Wrist(s)	360	1.2	14	10.6
33 Hand(s)- except finger(s)	40	0.1	6	29.6
34 Finger(s)- fingernail(s)	70	0.2	4	22.3
38 Multiple upper extremities locations	40	0.1	41	31.0
4 Lower extremities	530	1.8	13	9.1
41 Leg(s)	350	1.2	13	10.8

See footnotes at end of table

Table 2. Number, incidence rate ¹, median days away from work ² and relative standard errors ³ of occupational injuries and illnesses involving days away from work ⁴ to selected parts of body with musculoskeletal disorders⁵ in private industry for Georgia, 2007 -- Continued

Part of body affected	Total Cases	Incidence Rate	Median Days	Relative Standard Error
412 Knee(s)	310	1.1	15	11.2
42 Ankle(s)	100	0.4	9	18.6
43 Foot(feet)- except toe(s)	70	0.2	77	22.2
430 Foot(feet)- except toe(s)- unspecified	70	0.2	77	22.2
8 Multiple Body Parts	220	0.8	9	13.1

¹ Incidence rates represent the number of injuries and illnesses per 10,000 full-time workers and were calculated as: (N / EH) X 20,000,000 where,

N = number of injuries and illnesses,

EH = total hours worked by all employees during the calendar year,

20,000,000 = base for 10,000 full-time equivalent workers (working 40 hours per week, 50 weeks per year).

² Median days away from work is the measure used to summarize the varying lengths of absences from work among the cases with days away from work. Half the cases involved more days and half involved less days than a specified median. Median days away from work are represented in actual values.

³ Relative standard errors are a measure of the sampling error of an estimate. Sampling errors occur because observations are made on a sample, not on the entire population. Estimates based on the different possible samples of the same size and sample design could differ. Relative standard errors less than 0.05 are not shown.

⁴ Days away from work cases include those which result in days away from work with or without job transfer or restriction.

⁵ Includes cases where the nature of injury is: sprains, strains, tears; back pain, hurt back; soreness, pain, hurt, except back; carpal tunnel syndrome; hernia; or musculoskeletal system and connective tissue diseases and disorders and when the event or exposure leading to the injury or illness is: bodily reaction/bending, climbing, crawling, reaching, twisting; overexertion; or repetition. Cases of Raynaud's phenomenon, tarsal tunnel syndrome, and herniated spinal discs are not included. Although these cases may be considered MSD's, the survey classifies these cases in categories that also include non-MSD cases.

NOTE: Dashes indicate data that do not meet publication guidelines or data for incidence rates less than .05 per 10,000 full-time workers. The scientifically selected probability sample used was one of many possible samples, each of which could have produced different estimates. A measure of sampling variability for each estimate is available upon request.

SOURCE: Bureau of Labor Statistics, U.S. Department of Labor, February 04, 2010